

SG - 467

Total No. of Pages : 3

Seat No.	
----------	--

M.Sc. (Part - I) (Semester - II) Examination, March - 2023

CHEMISTRY (NEP-2020)

Analytical Chemistry-II

(Paper-VIII-CH-2.4, APCH 2.4, IND 2.4) (CBCS)

Sub. Code : 90166/90076

Day and Date : Tuesday, 20 - 06 - 2023

Total Marks: 80


Time : 10.30 a.m. to 01.30 p.m.

- Instructions:
- 1) Attempt in all the five questions.
  - 2) Question No.1 is compulsory.
  - 3) Attempt any Two questions from Section-I and any two from Section-II.
  - 4) Answers to all questions from should be writtem in one answer book.
  - 5) All questions carry equal marks.
  - 6) Figures to the right indicate full marks.

Q1) Answer the following questions:

[16]

- a) List the factors affecting chemical shift.
- b) Differentiate between protonated ion and adduct ion.
- c) How will you differentiate between organochlorine and organobromine compound using MS spectra.
- d) Calculate the wavelength ( $\lambda_{\max}$ ) of given examples 1,3,5-hexatriene



- e) How many signals does the aldehyde  $(\text{CH}_3)_3\text{CCH}_2\text{CHO}$  have in  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra?
- f) Define Beer's Lambert Law.

P.T.O.

- g) How will you differentiate trans-stilbene and cis-stilbene using UV-Vis spectroscopy?
- h) State the unique advantage of AAS in analysis of metals.
- i) What is thermogravimetric analysis?
- j) What is carrier gas in connection with thermal analysis and state its significance?
- k) What are the criteria for selecting the sample holders used in TG analysis?
- l) Compare between DSC and DTA.
- m) Which technique is generally used to deal with solid samples in AAS?
- n) Which of the following molecules will show IR spectrum  $\text{HCl}$ ,  $\text{CH}_4$ ,  $\text{CO}_2$ ,  $\text{H}_2$  and  $\text{N}_2\text{O}$ .
- o) How is sample prepared for AAS/ICP-AES?
- p) ICP-OES means

### SECTION - I

- Q2) a) What is spin-spin coupling? Explain with examples the types of spin-spin couplings. [8]
- b) Deduce the structure from the given data: [8]
- M.F. :  $\text{C}_8\text{H}_{11}\text{N}$ ; IR : 3350, 1596, 1020, 761, 703  $\text{cm}^{-1}$ .
- PMR:  $\delta$  7.2 (s, 5H), 3.87 q, 1H), 1.83 (s, 2H), 1.2 (d, 3H).
- $^{13}\text{C}$ -NMR: 148, 128, 126 125, 51, 26. Justify the spectral data.
- Q3) a) Explain the instrumentation, working and applications of mass spectrometry. [8]
- b) Explain the Woodward-Feiser rule for calculating absorption maxima with suitable examples. [4]
- c) State the applications of IR spectroscopy. How are primary, secondary and tertiary alcohols distinguished using IR spectroscopy? [4]



Q7. Write short note on **any four** of the following.

- State the advantages and disadvantages of AAS.
- Explain the shielding and deshielding effects.
- Compare between base peak and molecular ion peak
- Industrial Applications of DSC
- Different modes or types of thermogravimetric analysis.

### Spectroscopic Data

**Table: I** Some characteristic IR frequencies (Only approximate values)

$\equiv\text{CH}$ , 3300;  $=\text{CH}$ , 3050;  $\text{O}=\text{C}-\text{H}$ , 2800;  $\text{NH}$ , 3300;  $\text{O}-\text{H}$ , 3600;  $\text{C}\equiv\text{N}$ , 2200;  $\text{C}=\text{C}$ , 1620 to 1680; Aromatic, 1600-1500;  $\text{C}=\text{N}$ , 1660; Ketone, 1720; Ester, 1740; Saturated acids, 1720; Saturated aldehydes, 1730; Saturated amides, 1650;  $\text{CH}=\text{CH}_2$ ,  $\text{H}$  (CIS), 690;  $\text{C}=\text{H}$ , 790-840;  $\text{NO}_2$ , 1530 and 1350; Monosubstituted aromatics 690-710 and 730 to 770; Disubstituted 735-770; Trisubstituted 750-810; Tetra substituted, 770, 800-860;  $-\text{CO}-\text{CH}_2-\text{Cl}$ , 1745-1725 all values are in  $\text{cm}^{-1}$

**Table: II** Approximate chemical shifts of  $-\text{CH}_3$ ,  $-\text{CH}_2$  and  $-\text{CH}$  protons in  $\delta$  (ppm)

$\text{C}-\text{CH}_3$ , 0.9;  $\text{O}-\text{CH}_3$ , 1.6;  $\text{Ar}-\text{CH}_3$ , 2.2;  $\text{O}=\text{C}-\text{CH}_3$ , 2.3;  $\text{N}-\text{CH}_3$ , 2.3;  $\text{O}-\text{CH}_3$ , 3.3;  $\text{C}=\text{CH}_2$ , 4.6; to 5.3;  $\text{C}=\text{CH}$ ;  $\text{Ar}-\text{H}$ , 7 to 9;  $-\text{CO}-\text{C}=\text{CH}$ , 6.2;  $\text{C}-\text{CH}-\text{CO}$ , 5.7.

**Table: III** Approximate  $^{13}\text{C}$  chemical shifts

$\text{R}-\text{CH}_3$ , 530;  $\text{R}-\text{CH}_2-\text{R}$ , 25-55;  $\text{R}_3-\text{CH}$ , 35 to 70;  $\text{R}_4\text{C}$ , 30-50;  $\text{R}_3\text{C}-\text{O}$ , 57-80;  $\text{R}_3\text{C}-\text{N}$ , 60 to 75;  $\text{C}\equiv\text{C}$ , 75 to 105;  $\text{C}\equiv\text{N}$ , 110 to 125;  $\text{C}=\text{C}$ , 100 to 140; Aromatics 115 to 145;  $\text{R}-\text{COOR}$ / $\text{R}-\text{CONH}_2$ , 155-180;  $\text{R}-\text{COOH}$ , 165-185;  $\text{R}-\text{CHO}$ , 185-205;  $\text{RCOR}$ , 190-225

